



Bacteria Contamination and Distribution in Patient Care Environment at Mbagathi Hospital Surgical and New-born Units

Enock Ongeru Magori^{1*}, Stanley Kamwati Kang'ethe², Kennedy Kuria Muna³, Gunturu Revathi⁴ and Geoffrey Omuse⁴

¹Mount Kenya University/Aga Khan University P.O BOX 342-0100, Thika, Kenya

²Mount Kenya University P.O BOX 342-0100, Thika, Kenya

³Muranga University P.O BOX 75-0100, Muranga, Kenya

⁴Aga Khan University Hospital P.O BOX 30270-0100, Nairobi, Kenya

ABSTRACT

Background: Hospital environment can serve as an important reservoir and thus a critical element in the transmission of bacterial infections especially in critical care settings such as Surgical and new-born units. Contact with contaminated surfaces may lead to Hospital Acquired Infections (HAIs) among healthcare workers, visitors and patients. Further, HAIs may contribute to the spread of drug-resistant bacterial infections. This study sought to determine bacteriological characteristics and distribution in patient care environmental surfaces (Surgical and new-born units) at Mbagathi hospital Nairobi Kenya.

Methods: A total of 700 samples were obtained from different surfaces: beds, bedside tables, sink taps handle, door handles, nursery incubators, paediatric weighing scale and new-born resuscitation machine by means of repeated screens over a period of three months. Microbiological isolation and identification were done by following standard laboratory methods established at Aga Khan University Hospital. Antibiotic susceptibility testing was carried out by using Vitek 2 system.

Result: Five bacterial genera were isolated, *S. Aureus* was 3% (19/700), *E. coli* 0.9% (6/700), *Acinetobacter* species was 1.4% (10/700), *Pseudomonas* species 0.1% (1/700) and coagulase negative staphylococcus (CoNS) 13.0% (88/700). Trimethoprim/Sulfamethoxazole and Benzylpenicillin were the most resistant antimicrobial agents while Oxacillin, Cefoxitin, Levofloxacin, Linezolid was most sensitive.

Conclusion: All *S. aureus* were methicillin sensitive (MSSA). Patient beds surfaces were the most contaminated among the selected items while the nursery incubator was the least contaminated.

Keywords: Hospital Environment, Bacterial Contamination, Surfaces, Surgical, New-Born Unit

Introduction

Microorganisms are known to survive on inanimate surfaces for a long period^[1] and it is more important in healthcare setting environment where patients with immunodeficiency are at a higher risk of contracting hospital acquired infections. Fomites are items that become contaminated with pathogenic organisms and act as vectors in their transmission. Pathogens are shed from different body secretions during and after the course of infections. Fomites include patient care items and environment surfaces. Surface colonization of a variety of inanimate items in a hospital facility by microorganisms has been reported as a possible route for the transmission of many disease-causing organisms.^[2] Transmission of these pathogens is by direct or indirect contact with colonized or infected patients and contaminated instruments in the hospital.^[3] Mbagathi County Hospital is the largest referral hospital among the devolved units in Nairobi City County taking care of up to 800 inpatients every week with almost 2500 workers and likewise accepting at least 2300 guests

of patients every day. This populace makes it possible for contamination of highly touched areas.

The World Health Organization in 2014 acknowledged that antimicrobial resistance is a serious universal problematic area to public health. The most common multi resistant pathogens that have been documented are; *S. aureus*, *E. coli*, *P. aeruginosa*, *Enterococcus* species, *Acinetobacter* species, *K. pneumonia* which are frequently associated with skin and soft tissue infections.^[4] The same organisms have been isolated from fomites from hospital environment especially patient zone environment.^[5] Multidrug resistant organisms limit the curative options thus creating social and economic burden to healthcare system. These pathogens are responsible for disease, death and antimicrobial resistance in advanced and emerging developing countries.^[5] Medical microbiology laboratories in developing countries especially in Kenya are sparingly spread, poorly equipped because they are inadequately financed. These challenges the Medical laboratories encounter pose a great



danger and it is risk at this era of emergence of multidrug resistance organisms.

There is paucity of data on the prevalence of nosocomial infections and degree of bacterial contamination of mostly touched item surfaces in patient care environment and there are no records of the situation in Nairobi County mainly Mbagathi hospital. The previous studies that have been done on patients' clinical samples in chosen hospitals in Nairobi City County, MRSA was regularly isolated from public healthcare amenities attending economically underprivileged Nairobi populace such as those residing in metropolitan informal settlements.^[6] Other reports indicate there is transmission of MRSA and MSSA in public healthcare facilities in Kenya.^[7]

Materials and Methods

This study was conducted at Mbagathi County Hospital located in Dagoretti sub-county of Nairobi City County in Kenya between May and July 2020. The hospital has a 200 bed-capacity with approximately 800 admissions monthly and 1000 outpatient's visits per day where around 300 healthcare practitioners attend them. In general ward at Mbagathi Hospital, floors are mopped twice in a day by using detergent solutions and in corridors floors they are scrubbed by using mopping machines weekly. Working staff tables and benches are wiped once daily by using sodium hypochlorite solutions. There is no protocol of cleaning/ disinfecting surfaces of bed rails, nursery incubator, bed side table, water sink handle, ward door handle, paediatric weighing scale, ward drug trolley, new-born and resuscitation machine. The study was approved by Mount Kenya University, National Commission for Science, Technology and Innovation ethical committees. Authorization was also granted by the Mbagathi Hospital and Aga Khan University Hospital ethical committee.

Specimen Collection

A total of 700 samples were collected from different item surfaces within Surgical and New-born units. Samples were collected from bed rails, nursery incubator, bed side table, water sink handle, ward door handle, paediatric weighing scale, ward drug trolley, new-born and resuscitation machine. Samples from chosen surfaces were collected by rubbing and rotating sterile cotton swabs moistened with 0.9% normal saline firmly over the entire target surface which were conveniently selected from surgical and new-born units. These sites were more accessible and mostly used by patients, visitors and healthcare workers. These units were selected because of patient load and likelihood of long stay in the hospital.

Isolation and identification of bacterial isolates,

The swabs upon collection and well labelled, they were transported to Aga Khan University Hospital Laboratory in amies transport media. The specimens were inoculated within two hours after collection onto 5% sheep blood agar by streaking the agar surfaces with wire loop then incubated aerobically at 37 °C for up to 48 hours. After the end of incubation period, the culture plates were examined for bacterial growth and their morphological features of the organisms on the agar used. The identification of culture isolates was done according to standard operating procedures adapted at Aga Khan University Hospital. Conventional identification criteria included Gram staining, catalase, coagulase and oxidase tests. Further identification was performed by using the Vitek 2 system.

Antibiotic susceptibility test

An automated Antibiotic susceptibility to a standard panel of antibiotic was performed by Vitek 2 machine, version 4.01 (Biomerieux, Mercy-I' Etoile France) an automated system for bacterial identification which performs antibiotic susceptibility by using broth dilution. Also, Kirby disc diffusion method was used. Bacterial isolates that was resistant to three or more drug classes were categorized as multidrug resistant organisms (MDR)

Quality control

Culture media were incubated without inoculum to check the sterility of the culture media. Standard organisms were inoculated on 5%sheep blood agar and MacConkey to check their growth support. Standard organism used as control for identification and susceptibility are; *Staphylococcus aureus* (ATCC ID: 25923 and ATCC AST: 29213) *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853).

Result

Out of 700 collected samples from various item surfaces, growth was obtained in 63% (442/700) samples of which 18% (124/700) were identified as clinical pathogens. 37% (258/700) of the samples, there were no growth obtained. Various bacterial flora was isolated from different surfaces. The bacterial contamination in the beds, incubators, bedside tables and ward drug trolleys was 30.7%, 2.4%, 29.5% and 32% respectively.

Based on sample item sites, for bed, prevalence was highest in the right-hand side (48%) and lowest in the head and bottom sides (28%). Prevalence in incubators was highest in the top outside (12%). Prevalence was highest in the top (20%) in bedside tables, handle (20%) in ward doors and handle (10%) in water sinks.

The study found that five bacterial genera were isolated (*S. aureus*, *E. coli*, *Acinetobacter spp*, *Pseudomonas spp*, CoNS). It was determined that the prevalence of *S. aureus* bacterial contamination in patient care environment at Mbagathi hospital surgical and new-born unit was 3%, *E. coli* (0.9%), *Acinetobacter spp* (1.4%), *Pseudomonas spp* (0.1%), while the most dominant contaminant was CoNS (13%). Most *S. aureus* isolates were obtained from bed rails (12/19), ward door handles (3/19), drug trolley (1/19) and bedside table (3/19). All the *S. aureus* were MSSA as shown on table 4. Out of 19 *S. aureus* isolated, 11 were multidrug resistant (table 5). For Gram negative antimicrobial resistance pattern, it is demonstrated in table 4-7.

Discussion

The study found that item surfaces in surgical and New-born units that are commonly used by patients, visitors and Healthcare workers were contaminated with potential pathogens. The role of the items on how they are involved in infection transmission was not determined in this study. However, the isolation of *S. aureus*, Coagulase negative Staphylococci, *E. coli*, *Pseudomonas* species, *Acinetobacter* species makes it possible for the transmission of Hospital Acquired Infection. Nosocomial pathogens if there is no continuous disinfection of surfaces would be the constant source of transmission as these organisms are can survive on item surfaces for months.

Table 1: Bacterial distribution pattern on ward items surfaces.

	S. aureus	E. coli	Acinetobacter spp.	Pseudomonas spp.	CoNS	Total	P value
Surgical bed	12 (14.0)	5 (8.1)	7 (5.5)	1 (1.2)	61 (71.0)	86 (100.0)	
Incubator NBU	0 (0.0)	0 (0.0)	0(0.0)	0 (0.0)	2 (100.0)	2(100.0)	
Bedside table	3 (33.3)	1(11.1)	3(33.3)	0 (0.0)	2 (22.3)	9 (100.0)	.845
Water sink	0 (0.0)	0 (0.0)	0(0.0)	0 (0.0)	1 (100.0)	1 (100.0)	
Ward door	3 (75.0)	0 (0.0)	0 (0.0)	0(0.0)	1 (25.0)	4 (100.0)	
Ward trolley	1 (33.3)	0 (0.0)	0 (0.0)	0 (0.0)	2 (66.7)	3 (100.0)	

Table 2: Bacterial distribution pattern in ward item sites.

		S. aureus	E. coli	Acinetobacter spp.	Pseudomonas spp.	Cons	Total	P value
Bed sites	Head	2 (12.0)	2 (12.0)	0 (0.0)	0 (0.0)	13 (76.0)	17 (100.0)	.345
	Bottom	4 (20.0)	1 (5.0)	2 (10.0)	0 (0.0)	13 (65.0)	19 (100.0)	
	Left hand side	3 (12.0)	2 (8.0)	2(8.0)	0 (0.0)	18 (72.0)	25 (100.0)	
	Right hand side	4 (14.0)	0 (0.0)	3 (10.0)	1 (3.0)	21 (73.0)	29 (100.0)	
Incubator sites	Top outside	0 (0.0)	0 (0.0)	1(20.0)	0 (0.0)	4 (80.0)	5 (100.0)	-
Bedside table	Top	3(18.0)	1 (6.0)	2 (13.0)	0 (0.0)	10 (63.0)	16 (100.0)	-
	Handle	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	5 (100.0)	5 (100.0)	
Water sink	Handle	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (100.0)	2 (100.0)	-
Ward door	Handle	4(80.0)	0 (0.0)	0(0.0)	0 (0.0)	2 (20.0)	6 (100.0)	-

Table 3: Antimicrobial sensitivity patterns of isolated *S. aureus* stain. Multidrug resistant *S. Aureus* organism in patient care environment surfaces.

Antimicrobial agent	Resistant (N)	Sensitive (N)	Total (N)
Linezolid	0	19	19
Trimethoprim/Sulfamethoxazole	14	5	19
Benzympenicillin	14	5	19
Erythromycin	7	12	19
Clindamycin	4	14	19
Gentamicin	0	19	19

Antimicrobial agent	Resistant (N)	Sensitive (N)	Total (N)
Oxacillin	0	19	19
Cefoxitin	0	19	19
Levofloxacin	0	19	19
Tetracycline	6	13	19
Rifampicin	6	13	19

Table 4: Multidrug resistance nature of *S. Aureus* isolates.

Antimicrobial	No. Resistant
Trimethoprim/Sulfamethoxazole + Benzylpenicillin + Erythromycin + Clindamycin	4
Trimethoprim/Sulfamethoxazole + Benzylpenicillin + Rifampicin	4
Trimethoprim/Sulfamethoxazole + Benzylpenicillin + Tetracycline	3
Total	11

Table 5: Antimicrobial sensitivity patterns of isolated *E. coli*.

Antimicrobial agent	Resistant (N)	Sensitive (N)	Total (N)
Ciprofloxacin	2	4	6
Trimethoprim/Sulfamethoxazole	3	3	6
Ceftazidime	2	4	6
Gentamicin	2	4	6
Meropenem	1	5	6
Ceftriaxone	3	3	6

Table 6: Antimicrobial sensitivity patterns of isolated *Acinetobacter* species stains.

Antimicrobial agent	Resistant (N)	Sensitive (N)	Total (N)
Imipenem	1	9	10
Cefotaxime	6	4	10
Clotrimoxazole	5	5	10
Ciprofloxacin	4	6	10
Ceftazidime	6	4	10
Gentamicin	4	6	10

Table 7: Antimicrobial sensitivity patterns of isolated *Pseudomonas* species stains.

Antimicrobial agent	Resistant (N)	Sensitive (N)	Total (N)
Ceftriaxone	0	1	1
Trimethoprim/Sulfamethoxazole	1	0	1
Gentamicin	0	1	1
Ciprofloxacin	0	1	1
Meropenem	0	1	1
Ceftazidime	0	1	1

The findings when compared to other studies elsewhere, the prevalence rate in this study was 18% slightly lower than that reported by Nwankwo^[8] at Murtala Mohammed Specialist Hospital in Nigeria. The prevalence rate by Nwankwo was found to be 23.33% (95% CI= 20.28 -26.71%). However, Ensayef^[9] found the prevalence to be even lower at Al Imam Ali Hospital in Bagdad 4%. However, this prevalence was also lower than that reported by Al-Zoubi,^[10] whose prevalence was 98.7% at San Vicente Foundation, a teaching hospital in Columbia and Gondar University Hospital in which 83.1% (95% CI = 68.78 –99.52%) of inanimate objects were contaminated.

The study determined that the prevalence of *S. aureus* bacterial contamination in patient care environment at Mbagathi hospital surgical and new-born unit was 3%. This prevalence was lower than that recorded by Segujja^[11] who found the 167 (57.59%, CI = 49.18 - 67.01) harboured bacterial pathogens, where *S. aureus* prevalence was 25.75%. The prevalence of *E. coli* was 18.55%, a higher prevalence than that of Mbagathi hospital surgical and new-born unit, which was found to be 0.9%. However, the current study did not find existence of some bacteria such *E. faecalis*, *S. epidermidis*, *P. mirabilis*, *Bacillus* species and *S. saprophyticus* which were prevalent in Segujja^[11] study.

The study found that CoNS (13.0%) and *Staphylococcus aureus* (3%) were the most frequently isolated pathogen across the hospital surfaces, a consistent result with other findings from different studies and authors across the world.^[12,13] In this study, CoNS were the most prevalent in all item sites. *S. Aureus* were present in surgical beds, ward doors, bedside table and drug trolleys. The other type of bacteria did not cut across all surfaces, and only existed in one or two surfaces. In this regard, *E. coli* were only present in surgical beds. *Acinetobacter* species were present in surgical beds bedside table, nursery incubator. *Pseudomonas* species were only present in surgical beds.

The prevalence of *Acinetobacter* species at Mbagathi hospital surgical and new-born unit was 1.4%. In contrast to these findings, Carvalheira^[14] found that the most common species were *Acinetobacter guillouiae* with a prevalence of 34.9%. In addition, Banerjee *et al.* (2018) found a high prevalence of *Acinetobacter* species associated with HCAI in the adult ICU of a tertiary care hospital in Varanasi, north India. Moreover,^[15] also revealed that *Acinetobacter* species were the significant major pathogen against other Gram-negative organisms, which is the same finding to the present study. The prevalence of *Pseudomonas* species at Mbagathi hospital surgical and new-born unit was found to be 0.1%. These findings are different from those of

Banerjee^[15] who found the prevalence of *Pseudomonas spp* to be 10.17% in Intensive Care Unit of a Tertiary Care Hospital, Varanasi, India.

The prevalence of contamination in the beds, incubators, bedside tables and ward drug trolleys was 30.7%, 2.4%, 29.5% and 7% respectively. The prevalence of bacterial contamination was therefore highest in the beds. The findings when compared to other studies agree with those of Zazouli^[16] who found a significant prevalence of contamination in hospital beds. The overall contamination was 40%, which is slightly higher than that found in the beds in this study (36.5%). Hassan^[17] also found that the most regularly contaminated surfaces were the bed tops, and bed sides. These findings are also true as postulated in a study by Saka^[18].

Conclusion

From the findings obtained in this study, the study concludes that bacterial contamination at Mbagathi hospital surgical and new-born unit on item surfaces that are frequently used by healthcare workers, visitors and patients is 18%. Specifically, the prevalence of *S. aureus* was 3%, *E. coli* (0.9%), *Acinetobacter spp* (1.4%), *Pseudomonas spp* (0.1%), while the most dominant contamination was CoNS (13.0%). The isolation of pathogens from fomites demonstrates poor hand hygiene among visitors, healthcare workers and patients. These potential pathogens that were isolated from mostly touched surfaces signify the importance of establishing the cleaning protocol and disinfection of these sites. There is need to emphasize the importance of hand hygiene and decontamination of these sites regularly.

Acknowledgements

We thank Germaine research foundation for funding this study. We also thank microbiology staff of Aga Khan University Hospital and Mbagathi hospital for their technical support. We thank Mount Kenya University, Mbagathi and Aga Khan University Hospitals for giving us a chance to study and giving a chance to collect samples and carry out our experiments.

Funding

Germaine research foundation funded this project.

Competing Interests

None declared.

Reference

1. Russotto, V., Cortegiani, A., Raineri, S. M., Iozzo, P., Gregoretta, C., & Giarratano, A. (2017). What is the risk of acquiring bacteria from prior intensive care unit bed occupants?. *Critical Care*, 21(1), 55-62.

2. Olise, C. C., Simon-Oke, I. A., & Akumeh, D. D. (2018). Clinical isolates resistance to commonly used antibiotics: A concern in healthcare setting. *International Journey Public Health Safe*, 3(150), 2-4.
3. Facciola, A., Pellicano, G. F., Visalli, G., Paolucci, I. A., Rullo, E. V., Ceccarelli, M., ..., & La Fauci, V. (2019). The role of the hospital environment in the healthcare-associated infections: a general review of the literature. *European review for medical and pharmacological sciences*, 23(1), 1266-1278.
4. Pendleton, J. N., Gorman, S. P., & Gilmore, B. F. (2013). Clinical relevance of the ESKAPE pathogens. *Expert review of anti-infective therapy*, 11(3), 297-308.
5. Bhatta, D. R., Hamal, D., Shrestha, R., Subramanya, S. H., Baral, N., Singh, R. K., ... & Gokhale, S. (2018). Bacterial contamination of frequently touched objects in a tertiary care hospital of Pokhara, Nepal: how safe are our hands?. *Antimicrobial Resistance & Infection Control*, 7(1), 97.
6. Maina, E. K., Kiiyukia, C., Wamae, C. N., Waiyaki, P. G., & Kariuki, S. (2013). Characterization of methicillin-resistant *Staphylococcus aureus* from skin and soft tissue infections in patients in Nairobi, Kenya. *International journal of infectious diseases*, 17(2), e115-e119.
7. Oosterik, L. H., Peeters, L., Mutuku, I., Goddeeris, B. M., & Butaye, P. (2014). Susceptibility of avian pathogenic *Escherichia coli* from laying hens in Belgium to antibiotics and disinfectants and integron prevalence. *Avian diseases*, 58(2), 271-278.
8. Nwankwo, E. (2012). Isolation of pathogenic bacteria from fomites in the operating rooms of a specialist hospital in Kano, North-western Nigeria. *Pan African Medical Journal*, 12(1), 12-19.
9. Ensayef, S., Al Shalchi, S., & Sabbar, M. (2009). Microbial contamination in the operating theatre: a study in a hospital in Baghdad. *EMHJ-Eastern Mediterranean Health Journal*, 15(1), 219-223.
10. Al-Zoubi, M. S., Al-Tayyar, I. A., Hussein, E., Al Jabali, A., & Khudairat, S. (2015). Antimicrobial susceptibility pattern of *Staphylococcus aureus* isolated from clinical specimens in Northern area of Jordan. *Iranian journal of microbiology*, 7(5), 265-281.
11. Segujja, F., Mwambi, B., Drago, C. K., Lubowa, N. M., Mugambwa, J., & Wabuyi, P. (2016). Characterization and antimicrobial susceptibility patterns of isolates from ward fomites. Kampala, Uganda: Makerere University.
12. Feglo, P., & Afriyie-Asante, A. (2014). Environmental impact on postoperative wound infections in a privately owned hospital in Ghana. *African Journal of Microbiology Research*, 8(15), 1620-1626.
13. Yallem, W. W., Kumie, A., & Yehuala, F. M. (2016). Point prevalence of hospital-acquired infections in two teaching hospitals of Amhara region in Ethiopia. *Drug, healthcare and patient safety*, 8(1), 71-79.
14. Carvalheira, A., Casquete, R., Silva, J., & Teixeira, P. (2017). Prevalence and antimicrobial susceptibility of *Acinetobacter* spp. isolated from meat. *International journal of food microbiology*, 243(1), 58-63.
15. Banerjee, T., Mishra, A., Das, A., Sharma, S., Barman, H., & Yadav, G. (2018). High prevalence and endemicity of multidrug resistant *Acinetobacter* spp. in intensive care unit of a tertiary care Hospital, Varanasi, India. *Journal of pathogens*, 2018(1), 1-9.
16. Zazouli, M. A., Yazdani-charati, J., Ahanjan, M., & Eslamifard, M. (2017). Bacterial contamination of environmental surfaces in two educational hospitals under the auspices of Mazandaran University of Medical Sciences. *Journal of Health in the Field*, 3(1), 22-29.
17. Hassan, M. Z., Sturm-Ramirez, K., Rahman, M. Z., Hossain, K., Aleem, M. A., Bhuiyan, M. U., ... & Gurley, E. S. (2019). Contamination of hospital surfaces with respiratory pathogens in Bangladesh. *PloS one*, 14(10), e0224065.
18. Saka, K. H., Akanbi, A. A., Obasa, T. O., Raheem, R. A., & Oshodi, A. J. (2017). Bacterial Contamination of Hospital Surfaces According to Material Make, Last Time of Contact and Last Time of Cleaning/Disinfection. *Journal of Bacterial Parasitology*, 8(03), 8-11.

***Corresponding author:**

Enock Magori, Mount Kenya University/Aga Khan University P.O BOX 342-0100, Thika, Kenya

Phone: +254713489101

Email: Enockonger2010@gmail.com

Date of Submission : 04/12/2020

Date of Acceptance : 23/02/2021

Date of Publication : 30/03/2021

Financial or other Competing Interests: None.